



# Membrane excitability and excitation–contraction uncoupling in muscle fatigue

Michael Fauler<sup>a,\*</sup>, Karin Jurkat-Rott<sup>b</sup>, Frank Lehmann-Horn<sup>b</sup>

<sup>a</sup> Institute of Applied Physiology, Ulm University, Germany

<sup>b</sup> Division of Neurophysiology, Ulm University, Germany

## Abstract

High-frequency tetanic stimulation is associated with an increase in extracellular and T-tubular  $K^+$  and changes of  $Na^+$  and  $Cl^-$  concentrations, membrane depolarization as well as inactivation of voltage-gated  $Na^+$  channels. These alterations are expected to lead to fiber inexcitability, which is largely prevented by mechanisms intrinsic or extrinsic to muscle fibers. They act by adapting electrical membrane properties or by accelerating the reconstitution of ionic homeostasis. The high  $Cl^-$  conductance of muscle fibers supports the  $K^+$  conductance in fast and complete repolarization and creates a mechanism for the fast reuptake of  $K^+$ , thereby reducing the T-tubular  $K^+$  accumulation. Excitability is increased by a  $Ca^{2+}$  and protein kinase C dependent inhibition of the  $Cl^-$  conductance which is efficient especially in the T-tubular system. Several mediators activate the  $Na^+/K^+$ -ATPase and thus enhance the restoration of ionic homeostasis. Examples are purines (ATP, ADP), calcitonin-gene related peptide and adrenaline. It is also necessary to adapt the strength of the sarcoplasmic  $Ca^{2+}$  concentration to the requirements of tetanic contractions. An overwhelming  $Ca^{2+}$  signal leads to enzymatically driven excitation–contraction uncoupling. This process is most likely driven by the  $Ca^{2+}$  dependent protease  $\mu$ -calpain and might lead to the long-lasting fatigue observed after excessive physical activity.

© 2012 Elsevier B.V. All rights reserved.

**Keywords:** Exercise;  $Cl^-$  conductance; Extracellular  $K^+$ ; T-tubular system;  $Na^+$  channels;  $Na^+$ - $K^+$ -ATPase; Membrane potential

## 1. Introduction

The actuation of muscle fiber contraction is a complex process involving several distinct steps. It is initiated by the interplay of various brain regions and propagates over the spinal cord to primary motor neurons, which elicit an action potential at the neuromuscular junction. From there, an action potential wave has to spread along the sarcolemma to excite the complete surface of the muscle fiber. At openings of the transverse tubular system the excitation wave propagates deep into the fiber where electrical excitation is subsequently transduced into contraction by a process called excitation–contraction coupling. This is mainly

mediated by  $Ca^{2+}$  release from the sarcoplasmic reticulum (SR). At all stages of this complex process, beginning from the motivation to perform limb or body movement, across the initiation and coordination of stimulation of motor neurons, downstream to the muscle fiber itself and its subcellular domains, failure might occur which could be objectively expressed as weakness and might be perceived as fatigue [1]. Activation of a muscle fiber goes along with a sequence of distinct alterations in the physical state of cells followed by their recovery. Repetitive activation, as during exercise, can result in the prolonged persistence of an activated state with incomplete recovery between stimuli. If the rate of stimulation is higher than the rate of recovery, the accumulation of remainders can constitute a striking homeostatic challenge. An accumulation like the tetanic increase of sarcoplasmic  $Ca^{2+}$  concentration might be desirable or detrimental, since it increases force but might also trigger excitation–contraction (EC) uncoupling [2].

\* Corresponding author. Address: Institute of Applied Physiology, Ulm University, Albert-Einstein-Allee 11, 89081 Ulm, Germany. Tel.: +49 731 50036224.

E-mail address: [michael.fauler@uni-ulm.de](mailto:michael.fauler@uni-ulm.de) (M. Fauler).

Myogenic fatigue ultimately develops if the rate of muscle stimulation is higher than the rate of complete recovery. Homeostatic impairments occur at least on four different levels with different time scales: (i) electrical dis- and recharge of the membrane during the action potential, (ii) changes of  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Cl}^-$  concentrations, (iii)  $\text{Ca}^{2+}$  transients, and (iv) metabolic changes. The action potential represents a rapid discharge of the membrane capacity and its somewhat slower recharge. Since depolarizing currents are mediated by different charge carriers ( $\text{Na}^+$ ) than repolarizing currents ( $\text{K}^+$  and  $\text{Cl}^-$ ), every action potential is associated with changes of ion concentrations [3]. The complete restoration of ionic homeostasis is slower than the action potential itself and is not complete within the refractory period. Therefore, fast repetitive stimulation can represent a profound challenge to ionic homeostasis. It compromises excitability and needs mechanisms that upregulate  $\text{K}^+$  uptake and  $\text{Na}^+$  extrusion or an adaptive reduction of electrical membrane stability in order to avoid excitation failure. On a similar time-scale changes of free  $\text{Ca}^{2+}$  concentration occur.  $\text{Ca}^{2+}$  serves as a messenger that activates the contractile machinery but in addition has different signaling functions. The signal  $\text{Ca}^{2+}$  transmits (initiation of contraction or cell-signaling) depends on its spatial and temporal occurrence. A strong rise of the EC-coupling signal might be overwhelming and disrupt the microdomains of spatially and temporally organized information processing units within muscle fibers [2].

The contribution of reduced excitability to the development of muscle fatigue depends on the type, intensity and duration of exercise. Especially short bouts of high-intense muscle activity might lead to a decline of fiber excitability. More typical modes of exercise are affected by metabolic or other causes of fatigue. In such cases excitability is essentially maintained by several regulatory mechanisms within muscles. The preservation of excitability has to be traded off against the risk of  $\text{Ca}^{2+}$  overload which disturbs the physiological signaling to the contractile machinery and other  $\text{Ca}^{2+}$ -dependent signal proteins by the activation of proteases that uncouple the excitation–contraction interface [2]. Here we concentrate on electrical excitation of muscle fibers and excitation–contraction coupling in relation to the development of muscular fatigue during exercise. The paper is based on the comprehensive reviews by Ament et al. [1] and Allen et al. [3]. If not quoted otherwise, statements are reasoned by these reviews.

## 2. The balance between loss and maintenance of excitability

In respect of electrical excitation, a muscle fiber can be structurally and functionally separated into three distinct compartments: the neuromuscular junction, the sarcolemma and the T-tubular system. Each of these main domains is specialized to serve a distinct function. Their integrated cooperation ensures a reliable and precise electrical response to nerve stimulation. The neuromuscular junction is optimized to elicit a propagating wave of electri-

cal excitation on the sarcolemma with a high safety factor of impulse transmission. It is a highly differentiated structure with membrane infoldings to increase effective surface area and capacitance to provide a large current that excites the adjacent sarcolemma. Dysfunction typically leads to muscle weakness and fatigue in myasthenic syndromes. The sarcolemma longitudinally conducts the action potential wave with high velocity and activates T-tubules at their openings. The T-tubules direct the excitation wave deep into the muscle fiber and are the site where excitation–contraction coupling originates. The access resistance to the T-tubular system unloads the sarcolemmal current source from the very large T-tubular membrane capacitance that represents a big current sink. Thereby it is ensured that the sarcolemma is excited with high conduction velocity, but consequently the T-tubular membrane has to regenerate the action potential by its own equipment of voltage-gated ion channels [4], since pure electrotonic conduction would not be sufficient to activate the EC coupling machinery. Failure of proper excitation of the T-tubular system is typically seen e.g. in myotonia and paramyotonia congenita as well as periodic paralyses [5]. Changes of excitation properties of muscle fibers are quite common during exercise. Stimulation at high frequencies leads to a reduction of muscle fiber excitability at the sarcolemma and the T-tubular membrane which results in a reduction of conduction velocity or even block. In the surface electromyogram the amplitude and area of the muscular compound action potential is smaller and the spectral content is shifted to lower frequencies. It should also be mentioned that, due to the T-tubular access resistance, it is possible that the sarcolemma is well excitable while T-tubules fail [4]. In this case there won't be a reduction of the amplitude or area of the electromyogram.

Excitability can be defined as the amount of current (charge) needed to elicit an action potential, i.e. to reach the threshold for excitation. It is based on voltage- and time dependent properties of voltage-gated  $\text{Na}^+$  channels which, during an action potential, transit through a fixed sequence of conformational states (Fig. 1). At resting conditions about half of the channels are in a closed-activatable state. They open when the membrane potential is depolarized above the threshold of excitation, at which depolarizing currents (carried by  $\text{Na}^+$ ) are of the same size as hyperpolarizing currents (carried by  $\text{Cl}^-$  and  $\text{K}^+$ ). If the membrane is depolarized slightly above this threshold, depolarizing currents become continuously larger than hyperpolarizing currents, because voltage-gated  $\text{Na}^+$  channels open in a positive feedback fashion mediated by the depolarizing membrane potential. Time-dependent fast inactivation of voltage-gated  $\text{Na}^+$  channels stops the upstroke of the action potential. Repolarization of the membrane potential is mediated by an inward  $\text{Cl}^-$  flux and outward currents via voltage-gated  $\text{K}^+$  channels. Repolarization not only restores the membrane potential to its resting value but also the state of ion channels, e.g. voltage-gated  $\text{Na}^+$  and  $\text{Ca}^{2+}$  channels, which transit back




Fig. 1. High-frequency stimulation leads to weakness. Tetanic stimulation results in an increase in extracellular and T-tubular  $\text{K}^+$  concentration ( $[\text{K}^+]_o$ ) as well as a change of the electrochemical gradient of  $\text{Na}^+$  and consequently leads to membrane depolarization. This is followed by closed-state inactivation and incomplete recovery of voltage-gated  $\text{Na}^+$  channels ( $\text{Na}_V$ ) which will reduce excitability. State transitions of  $\text{Na}_V$  channels during an action potential are depicted in a highly schematic manner. The diagram (left panel) shows two action potentials. From a holding potential of  $-90$  mV a regular action potential is triggered (solid line) while from  $-60$  mV only a local non-conducted electrotonic deviation of the membrane potential is provoked (dashed line). Due to sarcolemmal excitation failure the effective size of motor units is reduced. Excitation failure of T-tubules will result in ineffective sarcomere activation. The  $\text{Cl}^-$ -conductance and  $\text{Na}^+\text{-K}^+\text{-ATPases}$  are able to delay excitation failure or even maintain excitability. Modified after [5].

from the inactivated to the closed-activatable state, a process that mainly determines the duration of the refractory period. The threshold of excitation is not a static property of a muscle fiber but a dynamically evolving property of every region of the cell membrane. It is determined by the fraction of voltage-gated  $\text{Na}^+$  channels that are in the closed-activatable state, by the conductance of  $\text{Cl}^-$  and  $\text{K}^+$  channels that mediate hyperpolarizing currents and thereby stabilize the resting membrane potential and by the strength and frequency spectrum of the stimulating current. The amount of activatable  $\text{Na}^+$  channels depends on the membrane potential. Depolarization by only a few millivolts reduces the fraction of activatable channels by a fast and direct transition into a closed-inactivated state and a slow inactivated state. A more pronounced depolarization into the sub-threshold region of membrane potentials (between  $-60$  and  $-50$  mV) is able to drastically impair the excitability of muscle fibers and could even lead to complete inexcitability and conduction block as in various types of periodic paralysis. But at least it causes a relevant prolongation of the refractory period such that the maximal stimulation frequency that can be followed is reduced (Fig. 1). During repetitive stimulation of muscle fibers, as in tetanic activation, there is a continuous decline of the fraction of activatable  $\text{Na}^+$  channels due to an incomplete

repolarization between action potentials based on changes in ionic homeostasis especially in the T-tubular system and  $\text{Na}^+$  channel (slow) inactivation. Changes of ion concentrations are a consequence of high-frequency stimulation since ion fluxes that occur during action potentials are not completely compensated in the short period between action potentials (Fig. 1). The discharge and subsequent recharge of the membrane capacity in the course of an action potential initially results in an exchange of  $\text{Na}^+$  for  $\text{K}^+$  leading to an increase or decrease of extracellular and T-tubular concentrations of  $\text{K}^+$  or  $\text{Na}^+$ , respectively. The restoration of  $\text{Na}^+$  and  $\text{K}^+$  distribution relies on the primarily active transport mediated by the  $\text{Na}^+\text{-K}^+\text{-ATPase}$  which is a relatively slow process. If the repolarization of the membrane potential had to rely completely on a 1:1 exchange of  $\text{Na}^+$  for  $\text{K}^+$ , changes of ion concentrations occurred very fast and would lead to membrane depolarization already after few action potentials. Especially the T-tubular system which has a very high surface to volume ratio is prone to such changes of ionic homeostasis. In muscle fibers a large part of the repolarization is carried by a  $\text{Cl}^-$  inward current. Thereby part of the necessary  $\text{Na}^+\text{-K}^+$  exchange is substituted by  $\text{Na}^+\text{-Cl}^-$  influx [6]. Subsequent changes of  $\text{Cl}^-$  concentrations result in a much slower change of the resting membrane potential. The

extracellular and especially T-tubular accumulation of  $K^+$  during activity depolarizes the equilibrium potential of  $K^+$  much faster than the equilibrium potential of  $Cl^-$  changes. The high resting  $Cl^-$  conductance (up to 8-fold higher than  $K^+$  conductance) pulls the membrane potential close to the  $Cl^-$  equilibrium potential. Therefore, it is possible that the membrane potential is hyperpolarized in relation to the  $K^+$  equilibrium potential, thus evoking a  $K^+$  inward current, e.g. via inward rectifying  $K^+$  channels. This mechanism assists in the clearance of  $K^+$  from the T-tubular system. But the ultimate restoration of ionic homeostasis still relies on  $Na^+-K^+$ -ATPases. Due to its high resting conductance, the chloride system also has impact on the threshold of excitation. This is of special importance for the T-tubular system, where stimulation occurs with relatively slow kinetics, because the electrical configuration of the T-tubular system in relation to the sarcolemma represents a low-pass filter [4]. High-frequency content of the propagating wave along the sarcolemma ensures the fast sarcolemmal conduction velocity, while the T-tubular system has to be activated primarily by the low-frequency content of the spreading wave.

The equilibrium potential for  $Cl^-$  settles down close to the time-averaged membrane potential by passive distribution of  $Cl^-$ . Every deviation from the resting membrane potential, e.g. by stimulation induces an opposing current that stabilizes the membrane potential.  $ClC-1$ , the predominant  $Cl^-$  channel in fibers of skeletal muscle, is an important site for regulation of excitability. During high-frequency stimulation  $Cl^-$  conductance is reduced in a first phase and, at least in fast-twitch muscle, later increases upon metabolic exhaustion [4]. The reduction of  $Cl^-$  conductance is mediated by a protein kinase C (PKC) dependent pathway. PKC is activated by  $Ca^{2+}$  released from the SR. In addition, a decrease of intracellular pH reduces  $Cl^-$  conductance, but during muscle activity the  $Ca^{2+}$ -PKC dependent pathway is more sensitive and has a higher gain. This actively regulated decrease of  $Cl^-$  conductance serves to maintain membrane excitability particularly in the T-tubular system [4]. It adapts  $Cl^-$  conductance to the reduced maximal voltage-gated  $Na^+$  current. SR released  $Ca^{2+}$  is a sufficient mediator of this response since a smoother action potential upstroke, a consequence of the reduced  $Na^+$  current, leads to a broadening of the action potential resulting in increased and prolonged sarcoplasmic  $Ca^{2+}$  elevations. Metabolic exhaustion results in a strong increase of membrane conductance in rat fast-twitch and probably also slow-twitch muscles [4]. This is due to the opening of ATP-sensitive  $K^+$  channels ( $K_{ATP}$ ) conductance increases about 14-fold) and an increase of  $Cl^-$  conductance (about 3-fold) by an unknown mechanism. The high resting conductance is able to block excitation. After cessation of stimulation recovery of initial resting conductance occurs within 1 min. The effect is sensitive to the extracellular glucose concentration. Another study demonstrated that ATP-sensitive  $K^+$  channels ( $K_{ATP}$ ) play a role in the smoothing of the sarcoplasmic  $Ca^{2+}$  concentration during

high-frequency stimulation [7]. Blocking of  $K_{ATP}$  channels with glibenclamide led to a very strong rise of the  $Ca^{2+}$  concentration and muscle force followed by a rapid decline within seconds below control conditions, thus representing increased fatigability. In addition  $Ca^{2+}$  concentration did not completely recover between short bouts (200 ms every second) of high-frequency stimulation, resulting in incomplete relaxation. Sarcolemmal and T-tubular membrane excitability is challenged by the aforementioned changes in ion concentrations, mainly the increase of extracellular or T-tubular  $K^+$  concentration. Although the chloride system is able to bridge over a rapid decay of excitability, it is pivotal that ionic homeostasis is restored. This can only be accomplished by the activity of  $Na^+-K^+$ -pumps. Hence, despite the intrinsic stimulating action of intracellular  $Na^+$ , a variety of auto-, para- and endocrine mediators exist that aim on the activation of  $Na^+-K^+$ -ATPases. Active muscle releases purines (ATP, ADP) possibly through pannexin-1 hemichannels [8]. Extracellular ATP stimulates ionotropic (P2X) and metabotropic (P2Y) receptors on the sarcolemma leading to the expression of myokines (IL6) and the stimulation of the  $Na^+-K^+$ -ATPase. Changes of  $Na^+-K^+$ -ATPase  $Na^+$  affinity and phosphorylation of phospholemman, a  $Na^+-K^+$ -pump inhibitor that is repressed if phosphorylated, are mediated by a P2Y-independent pathway [9]. Calcitonin-gene related peptide (CGRP) has been shown to stimulate  $Na^+-K^+$ -ATPases in skeletal muscle, but its fatigue-ameliorating effects are mainly mediated by another largely unknown mechanism. Probably it acts by shifting the voltage-dependence of slow inactivation of voltage-gated  $Na^+$  channels [10]. This would increase the number of activatable channels and thereby increase excitability. CGRP is released together with acetylcholine at the neuromuscular junction and has been suggested to be responsible for the walk-through phenomenon seen in patients with hyperkalemic periodic paralysis [11]. Adrenaline is a strong stimulator of  $Na^+-K^+$ -ATPases. The effect is mediated by a protein kinase A dependent pathway via  $\beta_2$ -adrenergic receptors. It is able to improve excitability at high-frequency stimulation. Similar results have been achieved with insulin [3]. These results are derived from in vitro experiments on muscle preparations from rat. Studies employing healthy human volunteers on effects of the  $\beta_2$ -agonist salbutamol on fatigability did not show any significant improvement [12], neither did oral  $\beta$ -blockade has a measurable effect on excitability or myogenic fatigability but slowed the recovery of  $K^+$  homeostasis [13].

### 3. Excitation–contraction uncoupling in muscle fatigue

The breakdown of ATP during exercise leads to an accumulation of inorganic phosphate ( $P_i$ ) in the cytoplasm and the release of  $Mg^{2+}$  from its complex with ATP [1].  $Mg^{2+}$  is able to counteract the release of  $Ca^{2+}$  from the SR.  $P_i$  also inhibits  $Ca^{2+}$  release by building  $Ca^{2+}-P_i$  complexes within the SR and by phosphorylating ryanodine receptors

(RyRs), the  $\text{Ca}^{2+}$  release channels of the SR. These mechanisms reduce the  $\text{Ca}^{2+}$  transient caused by electrical excitation and therefore represent a form of functional EC uncoupling. EC uncoupling in a narrower sense is a phenomenon based on the ultrastructural disruption of the molecular interface between the voltage-sensor L-type  $\text{Ca}^{2+}$  channel in the T-tubular membrane and the SR  $\text{Ca}^{2+}$  release channel (RyR) [2]. This complex of proteins is localized to the triad junction and is the molecular correlate of the EC coupling process (Fig. 2). The detailed molecular assembly of the junction is still unknown. Disruption of this interaction results in a strong decrease of  $\text{Ca}^{2+}$  release and force development upon electrical excitation, since  $\text{Ca}^{2+}$  influx through L-type  $\text{Ca}^{2+}$  channels of skeletal muscle is quite low and  $\text{Ca}^{2+}$  induced  $\text{Ca}^{2+}$  release too weak to trigger a regular response. EC uncoupling is believed to be responsible for long-lasting fatigue as it occurs after eccentric or excessive muscle activation and the weakness seen in some types of muscular dystrophies and maybe other myopathies [2]. The disruption of the molecular interface between L-type  $\text{Ca}^{2+}$  channels and RyR is of enzymatic nature. It is likely that it is accomplished by the  $\text{Ca}^{2+}$ -dependent cysteine protease  $\mu$ -calpain, an assumption based on the subcellular localization of this protease to the triad junction in uncoupled fibers, various biochemical and pharmacological properties of the uncoupling process and the fact that exogenously applied active  $\mu$ -calpain causes partial uncoupling of muscle fibers [2]. EC uncoupling is triggered by a strong sarcoplasmic  $\text{Ca}^{2+}$  load which can be achieved either by an exceptional high  $\text{Ca}^{2+}$  peak during tetanic contraction lasting for seconds or a prolonged  $\text{Ca}^{2+}$  signal at a relatively high level. It is interesting in this context that, in the study mentioned in the previous section which investigated the role of  $\text{K}_{\text{ATP}}$  channels in repetitive bouts of fatigue, muscles in which  $\text{K}_{\text{ATP}}$  channels were either inhibited or which stem from a  $\text{K}_{\text{ATP}}$  knock-out mouse showed a delayed and less complete recovery from fatigue [7]. In these muscles sarcoplas-

mic peak  $\text{Ca}^{2+}$  concentration was doubled and did not completely decline between tetanic stimulations. It is further interesting that a second bout of fatigue applied at least 30 min after the first seemed to be independent from  $\text{K}_{\text{ATP}}$  channels and showed a similar time-course of cytoplasmic  $\text{Ca}^{2+}$  and force development as under control conditions during the first bout. The authors called this phenomenon “fatigue pre-conditioning” and differentiated it from ischemic pre-conditioning by showing its independence from adenosine receptor signaling, inhibiting mitochondrial  $\text{K}_{\text{ATP}}$  channels and PKC as well as the use of scavengers of reactive oxygen species. If fatigue pre-conditioning were based on EC uncoupling [14], it would give some functional significance to EC uncoupling in respect to improving muscle performance by shaping the time-course of the cytoplasmic  $\text{Ca}^{2+}$  signal. In heart muscle it has been suggested that calpain-dependent cleavage of an auto-inhibitory domain in the serine/threonine phosphatase calcineurin leads to the activation of the transcription factor NFAT (nuclear factor of activated T-cells) which triggers a genetic program resulting in myocardial hypertrophy [15]. In skeletal muscle NFAT has been associated with the switch to a slow-twitch fatigue-resistant fiber phenotype [16].

#### 4. Concluding remarks

Muscle fatigue evolves due to the slow kinetics of complete recovery from single twitches which is leading to the accumulation of remainders during repetitive stimulation. With regard to sarcolemmal and T-tubular membrane excitability this is expressed by a continuous increment of inactivated voltage-gated  $\text{Na}^+$  channels, the extracellular and T-tubular accumulation of  $\text{K}^+$  and changes of  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations as well as membrane depolarization. These changes synergistically challenge excitability and are expected, especially at high-frequency stimulation, to contribute to muscle fatigue. In healthy subjects this seems to be avoided by several muscle intrinsic and extrinsic mechanisms that act to maintain excitability. Muscle intrinsic processes involve the high  $\text{Cl}^-$  conductance which partly relieves the  $\text{K}^+$  system from its burden to repolarize especially the T-tubular membrane and that assists  $\text{Na}^+$ - $\text{K}^+$ -ATPases in the clearance of  $\text{K}^+$  by hyperpolarizing the membrane potential beyond the equilibrium potential of  $\text{K}^+$ , thereby reducing the T-tubular and extracellular increase of  $\text{K}^+$  concentrations. Activation of PKC by cytoplasmic  $\text{Ca}^{2+}$  leads to an inhibition of  $\text{Cl}^-$  channels, thus probably adapting resting membrane conductance to a reduced availability of voltage-gated  $\text{Na}^+$  channels and decreased  $\text{Na}^+$  current. The  $\text{Na}^+$ - $\text{K}^+$ -ATPase is activated during exercise to accelerate the recovery of ionic homeostasis. This is intrinsically accomplished by an increased intracellular  $\text{Na}^+$  concentration but also by auto-, para- and endocrine mediators. Purines like ATP and ADP released from active muscle stimulate  $\text{Na}^+$ - $\text{K}^+$ -ATPases. CGRP is released at the neuromuscular junction together




Fig. 2. Mechanism of excitation–contraction (EC) uncoupling. The  $\text{Ca}^{2+}$ -dependent cysteine protease  $\mu$ -calpain is believed to disrupt the molecular interaction site between the voltage-sensor of the T-tubular membrane (dihydropyridine receptor, DHPR) and the  $\text{Ca}^{2+}$ -release channel of the sarcoplasmic reticulum (ryanodine receptor, RyR1). Modified after [5].

with acetylcholine and has been shown to activate  $\text{Na}^+$ - $\text{K}^+$ -ATPases, but probably does also affect the inactivation kinetics of voltage-gated  $\text{Na}^+$  channels. Adrenaline is released during exercise. By activating  $\beta_2$ -adrenergic receptors it increases  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity. Although not efficient in healthy individuals, the  $\beta_2$ -agonist salbutamol improves performance in hyperkalemic periodic paralysis and this might also hold for other muscle diseases in which T-tubular  $\text{K}^+$  accumulation is a relevant factor for pathogenesis or in which  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity is downregulated, e.g. in various myopathies like muscular dystrophies or McArdle disease. The content of  $\text{Na}^+$ - $\text{K}^+$ -pumps in muscle depends on an individual's activity: training leads to higher while inactivity reduces  $\text{Na}^+$ - $\text{K}^+$ -pump expression. This underscores the importance of training in all kinds of diseases that are linked to reduced physical activity. Among the many changes physical inactivity provokes in the musculature, the decline of its ability to restore  $\text{K}^+$  homeostasis is of general importance. In muscular diseases the sequence from sparse physical activity, over muscular remodeling followed by early exhaustion and increased fatigability, leading to insufficient activity could end in a vicious cycle. Despite para- and endocrine mediators other extrinsic factors contribute to the maintenance of excitability during muscle activity. Of special importance might be the "wisdom" of the central nervous system to switch between motor units before fatigue arises and to adapt, i.e. reduce the rate of motor neuron discharge to the prolonged refractory period of muscle fibers – the so called "muscle wisdom" hypothesis. EC uncoupling seems to be an important mechanism for the causation of long-lasting muscle fatigue. It is invoked by overwhelming  $\text{Ca}^{2+}$  concentrations that probably activate the cysteine protease  $\mu$ -calpain which is localized to the triad junction and cleaves the molecular bridge between the voltage-sensor and the  $\text{Ca}^{2+}$  release channel of the excitation–contraction coupling machinery. This procedure might protect muscle fibers from toxic side-effects of prolonged and high cytoplasmic  $\text{Ca}^{2+}$  concentrations and assist in the proper shaping of the  $\text{Ca}^{2+}$  signal for optimal tetanic contraction. In addition, it might produce a signal to cell nuclei which induces muscle plasticity by a mechanism similar to what has been proposed for cardiac myocytes.

## 5. Conflict of interest

None.

## Acknowledgement

The work was supported by research Grants from the Else Kröner-Fresenius-Stiftung (2010\_A27) and the BMBF (IonoNeuroNet). Frank Lehmann-Horn is endowed Senior Research Professor of the non-profit Hertie-Foundation.

## References

- [1] Ament W, Verkerke GJ. Exercise and fatigue. *Sports Med* 2009;39:389–422.
- [2] Lamb GD. Mechanisms of excitation–contraction uncoupling relevant to activity-induced muscle fatigue. *Appl Physiol Nutr Metab* 2009;34:368–72.
- [3] Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev* 2008;88:287–332.
- [4] Pedersen TH, Huang CL-H, Fraser JA. An analysis of the relationships between subthreshold electrical properties and excitability in skeletal muscle. *J Gen Physiol* 2011;138:73–93.
- [5] Lehmann-Horn F, Jurkat-Rott K. Voltage-gated ion channels and hereditary disease. *Physiol Rev* 1999;79:1317–72.
- [6] Dutka TL, Murphy RM, Stephenson DG, Lamb GD. Chloride conductance in the transverse tubular system of rat skeletal muscle fibres: importance in excitation–contraction coupling and fatigue. *J Physiol* 2008;586:875–87.
- [7] Boudreault L, Cifelli C, Bourassa F, Scott K, Renaud JM. Fatigue preconditioning increases fatigue resistance in mouse flexor digitorum brevis muscles with non-functioning K(ATP) channels. *J Physiol* 2010;588:4549–62.
- [8] Buvinic S, Almarza G, Bustamante M, et al. ATP released by electrical stimuli elicits calcium transients and gene expression in skeletal muscle. *J Biol Chem* 2009;284:34490–505.
- [9] Walas H, Juel C. Purinergic activation of rat skeletal muscle membranes increases  $V_{\text{max}}$  and  $\text{Na}^+$  affinity of the Na, K-ATPase and phosphorylates phospholemman and alpha1 subunits. *J Biol Chem* 2012;463:319–26.
- [10] Macdonald WA, Nielsen OB, Clausen T. Effects of calcitonin gene-related peptide on rat soleus muscle excitability: mechanisms and physiological significance. *Am J Physiol Regul Integr Comp Physiol* 2008;295:R1214–23.
- [11] Clausen T, Nielsen OB, Clausen JD, Pedersen TH, Hayward LJ.  $\text{Na}^+$ ,  $\text{K}^+$ -pump stimulation improves contractility in isolated muscles of mice with hyperkalemic periodic paralysis. *J Gen Physiol* 2011;138:117–30.
- [12] Crivelli G, Millet GP, Gremion G, Borrani F. Effects of salbutamol on the contractile properties of human skeletal muscle before and after fatigue. *Acta Physiol (Oxf)* 2011;203:311–20.
- [13] Unsworth K, Hicks A, McKelvie R. The effect of beta-blockade on plasma potassium concentrations and muscle excitability following static exercise. *Pflugers Arch* 1998;436:449–56.
- [14] Bruton J. What does the membrane K(ATP) channel really do in skeletal muscle? *J Physiol* 2010;588:4333.
- [15] Heidrich FM, Ehrlich BE. Calcium, calpains, and cardiac hypertrophy: a new link. *Circ Res* 2009;104:e19–20.
- [16] Gundersen K. Excitation–transcription coupling in skeletal muscle: the molecular pathways of exercise. *Biol Rev Camb Philos Soc* 2011;86:564–600.