

SCN4A-associated hypokalemic periodic paralysis merits a trial of acetazolamide

S.L. Venance, MD, PhD; K. Jurkat-Rott, MD;
F. Lehmann-Horn, MD, PhD; and R. Tawil, MD

Primary hypokalemic periodic paralysis (HypoPP) is characterized by episodes of transient flaccid paralysis in association with reduced serum potassium levels.¹ Onset of transient weakness is typically in the first or second decade, often on awakening. Weakness may be mild and focal or progress to flaccid quadriplegia, and the duration may last hours to days. Interindividual attack frequency varies, from daily episodes to a single lifetime episode, commonly decreasing with age. Predictable triggers are rest after exercise and high carbohydrate meals. Unlike hyperkalemic periodic paralysis, myotonia is absent. A proportion of affected individuals have slowly progressive permanent weakness.

Inheritance is autosomal dominant, although one-third of cases are sporadic, and penetrance is reduced in women. HypoPP is genetically heterogeneous. Three common mutations in *CACNA1S*, the α -subunit of the skeletal muscle L-type calcium channel gene,² account for 70% of cases (HypoPP1), whereas 10%³ of patients have a mutation in *SCN4A*, the α -subunit of the voltage-gated sodium channel gene (HypoPP2),^{4,5} and are clinically indistinguishable.

Patients with HypoPP are treated with administration of oral potassium at the time of an attack. The carbonic anhydrase inhibitors acetazolamide (ACZ)¹ and dichlorphenamide, as well as potassium-sparing diuretics, are used for prophylaxis. In some patients, ACZ results in exacerbation of symptoms,^{3,6} a trait that runs true in families.⁶ Two kindreds carrying mutations in the *SCN4A* gene (HypoPP2)^{3,6} led some researchers to postulate that all HypoPP2 patients might react adversely to carbonic anhydrase inhibitors.³ Here we report a HypoPP2 patient treated with ACZ without exacerbation of his episodic attacks of weakness.

Case report. A 29-year-old man had attacks of transient weakness since age 15 years. His family history was negative. Ictal potassium levels were reduced (2.5 to 2.9 mmol/L). Provocative triggers included rest after exercise, carbohydrate or alcohol intake, diarrheal illness, and poor sleep. Respiratory muscles were not involved, and there was no interictal weakness. Interictal neurologic examination was normal, and he had no clinical or electrical myotonia. An exercise test (of abductor digiti minimi) conducted off-treatment showed an increment in amplitude (17%) and area (89%) at 2 minutes postexercise and a maximal decrement in amplitude (38%) and area (59%) at 30 minutes postexercise. DNA testing revealed an R672S mutation in the *SCN4A* gene (Ulm, Germany).

He successfully managed his periodic paralysis with potassium supplementation at the beginning of attacks and before engaging in vigorous physical activity and took ACZ (250 mg/d) for prophylaxis. Episodes of weakness were initially characterized as mild (not interfering with activity, occurring daily, and lasting minutes), moderate (interfering with but not preventing activity, occurring three times a week, with duration of 1 hour), and severe (unable to function, occurring once a month, with duration up to 6 hours). Although never attack free for >1 week, episodes are less frequent with his present treatment regimen (mild, daily; moderate, once weekly; and severe, quarterly).

Discussion. ACZ is used for prevention of attacks in periodic paralysis.¹ However, recent reports of ACZ-induced worsening of symptoms in patients with HypoPP2 led to suggestions that ACZ should be avoided in such patients. The patient with HypoPP2 described here, with an R672S *SCN4A* mutation, was successfully treated with ACZ without aggravation of attacks. Similarly, the proband in a kindred with the R672S mutation in *SCN4A* had

Table Patients with *SCN4A* mutations: Response to acetazolamide (ACE) treatment

Reference	No. of patients	Sex	Mutation	ACZ treatment
Bulman et al., 1999	2 (1 family)	M	R669H	Better
Bendahhou et al., 2001	3 (1 family)	M	R672S	Worsened
Davies et al., 2001	1 (sporadic)	M	R672S	Better
Sternberg et al., 2001	3 (1 family)	M	R672G	Worsened
Venance et al., 2004	1 (sporadic)	M	R672S	Better

decreased attack frequency when treated with ACZ 250 mg twice daily and spironolactone 100 mg daily.⁷

The variability in responsiveness apparent with ACZ treatment in individuals with missense mutations in the sodium channel (table) suggests the modifying influence of other unidentified genetic and environmental factors. ACZ and other carbonic anhydrase inhibitors should continue to be considered for patients with HypoPP2 as a treatment option along with other potassium-sparing diuretics.

From the Neuromuscular Disease Center (Drs. Venance and Tawil), Department of Neurology, University of Rochester School of Medicine and Dentistry, NY; and the Department of Applied Physiology (Drs. Jurkat-Rott and Lehmann-Horn), Ulm University, Ulm, Germany.

Supported by a CINCH training fellowship under NIH U1 U54 RR019482-01 (S.L.V.) and by the Muscular Dystrophy Association.

Received April 26, 2004. Accepted in final form June 8, 2004.

Address correspondence and reprint requests to Dr. R. Tawil, Neuromuscular Disease Center, Department of Neurology, University of Rochester School of Medicine and Dentistry, 601 Elmwood Avenue, Box 673, Rochester, NY 14642-8673; e-mail: rabi_tawil@urmc.rochester.edu

Copyright © 2004 by AAN Enterprises, Inc.

References

- Resnick JS, Engel WK, Griggs RC, Stam AC. Acetazolamide prophylaxis in hypokalemic periodic paralysis. *N Engl J Med* 1968;278:582–586.
- Jurkat-Rott K, Lehmann-Horn F, Elbaz A, et al. A calcium channel mutation causing hypokalemic periodic paralysis. *Hum Mol Genet* 1994; 3:1415–1419.
- Sternberg D, Maisonobe T, Jurkat-Rott K, et al. Hypokalaemic periodic paralysis type 2 caused by mutations at codon 672 in the muscle sodium channel gene *SCN4A*. *Brain* 2001;124:1091–1099.
- Bulman DE, Scoggan KA, van Oene MD, et al. A novel sodium channel mutation in a family with hypokalemic periodic paralysis. *Neurology* 1999;53:1932–1936.
- Jurkat-Rott K, Mitrovic N, Hang C, et al. Voltage sensor sodium channel mutations cause hypokalemic periodic paralysis type 2 by enhanced inactivation and reduced current. *Proc Natl Acad Sci USA* 2000;97:9549–9554.
- Bendahhou S, Cummins TR, Griggs RC, Fu Y-H, Ptáček LJ. Sodium channel inactivation defects are associated with acetazolamide-exacerbated hypokalemic periodic paralysis. *Ann Neurol* 2001;50:417–420.
- Davies NP, Eunson LH, Samuel M, Hanna MG. Sodium channel gene mutations in hypokalemic periodic paralysis: an uncommon cause in the UK. *Neurology* 2001;57:1323–1325.

Neurology[®]

SCN4A-associated hypokalemic periodic paralysis merits a trial of acetazolamide

S. L. Venance, K. Jurkat-Rott, F. Lehmann-Horn, et al.

Neurology 2004;63;1977

DOI 10.1212/01.WNL.0000143068.99794.5B

This information is current as of November 22, 2004

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/63/10/1977.full
References	This article cites 6 articles, 3 of which you can access for free at: http://n.neurology.org/content/63/10/1977.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/63/10/1977.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical trials http://n.neurology.org/cgi/collection/all_clinical_trials Chromosomes http://n.neurology.org/cgi/collection/chromosomes Clinical trials Observational study (Cohort, Case control) http://n.neurology.org/cgi/collection/clinical_trials_observational_study_cohort_case_control Muscle disease http://n.neurology.org/cgi/collection/muscle_disease
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

