

Case Report/Case Series

Hypokalemic Periodic Paralysis Induced by Thymic Hyperplasia and Relieved by Thymectomy

Renrong Yang, MD; Karin Jurkat-Rott, MD; Jinlin Cao, MD; Guofeng Wang, MD; Hans-Peter Seelig, MD; Changping Yang, MD; Guibao Liu, MD; Lin Pan, MD; Haiyan Zheng, MD; Frank Lehmann-Horn, MD

IMPORTANCE Hypokalemic periodic paralysis is a muscle channelopathy based on mutations or predisposing variants or secondary to potassium wasting. In contrast to myasthenia gravis, an association with thymic hyperplasia has not yet been reported, to our knowledge.

OBSERVATIONS We report a male patient in his mid-20s with progressive episodes of flaccid muscle weakness, associated low serum potassium levels, and a pathologic decrement in the long exercise test. Because the familial inheritance in the family was initially unknown, thorough diagnostic tests were performed including contrast-enhanced computed tomography scan, which displayed a mass in the anterior mediastinum. The test results for autoantibodies against myasthenia gravis (acetylcholine receptor, muscle-specific tyrosine kinase, and low-density lipoprotein receptor-related protein 4) and other end plate channelopathies were negative, and test results for hypokalemia-inducing hormones (thyroid, corticotropin, and cortisol) were negative. Surgery identified a thymus of $13 \times 8 \times 3$ cm³. Histologic analysis was consistent with thymic hyperplasia of the follicular subtype and immunohistologic analysis showed cytokeratin 5/6 in hyperplastic epithelial cells. A 2-year follow-up revealed the postoperative absence of weakness episodes. As in 30% of familial cases, molecular genetics testing failed to identify a mutation in periodic paralysis genes.

CONCLUSIONS AND RELEVANCE Thymic hyperplasia can clinically manifest susceptibility to hypokalemic periodic paralysis. For patients with late onset or increasing weakness episodes, we recommend imaging to assess for thymic enlargement and thymectomy at thymic hyperplasia.

JAMA Neurol. 2013;70(11):1436-1439. doi:10.1001/jamaneurol.2013.3918
Published online September 23, 2013.

Author Affiliations: Department of Thoracic Surgery, 117 PLA Hospital, Hangzhou, China (R. Yang, Cao, C. Yang, Liu, Zheng); Division of Neurophysiology, Neuromuscular Disease Center, and Rare Disease Center, Ulm University, Ulm, Germany (Jurkat-Rott, Lehmann-Horn); Department of Pathology, Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou, China (Wang); Medizinisches Versorgungszentrum, Karlsruhe, Germany (Seelig); Department of Neurology, 117 PLA Hospital, Hangzhou, China (Pan).

Corresponding Author: Frank Lehmann-Horn, MD, Division of Neurophysiology, Ulm University, Albert-Einstein-Allee 11, 89081 Ulm, Germany (frank.lehmann-horn@uni-ulm.de).

Three types of hypokalemic periodic paralysis (PP) are known¹: (1) A dominant type due to disease-causing mutations of genes *CACNA1S*, *SCN4A*, and *KCNJ2* encoding sarcolemmal ion channels²⁻⁴; the mutations increase the probability of membrane depolarization at low serum potassium levels, thereby rendering the muscle fibers unexcitable.⁵ Disease prevalence is 1:100 000 and onset of weakness episodes usually is puberty.¹ (2) The “sporadic” type is based on variants of susceptibility genes. Susceptibility genes are *KCNJ18*, a channel gene located on chromosome 17q24.3, and perhaps additional genes on this locus encoding sarcolemmal ion channels.⁶⁻⁸ Clinical manifestation requires hyperthyroidism or Graves disease. This coincidence leads to the most common PP, occurring in 13% to 24% of Asian men with hyperthyroidism. (3) The third type is caused by chronic hypokalemia due to hereditary or sporadic potassium wastage, such as Conn or Bartter syndromes, licorice abuse, and an overdose of potassium-wasting diuretics.¹

In the 3 types, the clinical presentation is similar. A weakness episode is triggered by carbohydrate- and sodium-rich foods, infections, elevated cortisol levels (emotional stress and operation), and rest following physical stress. The disposi-

tion to sarcolemmal depolarization is demonstrated by a decrement in the long exercise test.¹ Permanent weakness in addition to episodes can occur.⁵ Ictal potassium ingestion and long-term administration of carbonic anhydrase inhibitors or aldosterone antagonists often improve the episodic or permanent weakness, respectively.⁵ Because hyperthyroidism and potassium wastage can be cured, careful screening for these conditions is mandatory.


For the first time, to our knowledge, we describe a patient whose hypokalemic weakness attacks were induced by thymic hyperplasia and abolished by thymectomy. The possibility of hypokalemic PP due to autoimmune channelopathy as a fourth type is discussed.

Report of a Case

Methods

Members of the family participating in the study gave their informed consent. Experiments were approved by the ethics committees of 117 PLA Hospital Hangzhou and Ulm Univer-

Figure 1. Long Exercise Tests



The compound muscle action potentials of the patient and his affected uncle show a typical late decrement in contrast to the healthy control. For details, see the Methods subsection.

sity. Chest imaging was performed by radiography and contrast-enhanced computed tomography (CT). The following serum concentrations were determined: electrolytes; corticotropin (ACTH); cortisol; aldosterone; thyroid hormones; creatine kinase; IgA, IgG, IgM, and IgE; light chains κ and λ , κ/λ ; complements C3 and C4; CD3, CD4, CD8, CD19, and CD4/CD8; and natural killer cells. The following autoantibodies were tested: acetylcholine receptor, muscle-specific tyrosine kinase, N- and T-type calcium channels, voltage-gated potassium channel complex, N-methyl-D-aspartate subunit R1, AMPA1/2, γ -aminobutyric acid B receptor, CASPR2, LGI1, RyR1, titin, skm, nuclear, ribonuclear, Sm, Sjögren syndrome A and B, PM-Scl, jo-1, centromere protein B, proliferating cell nucleus, DNA, nucleosome, histone, ribosomal proteins, and mitochondria. The recently reported myasthenia gravis autoantibody against the low-density lipoprotein receptor-related protein 4 (Lrp4), which functions as a muscle-specific tyrosine kinase ligand,^{9,10} was determined using an indirect immunofluorescence test on HEK293 cells, transiently transfected for 48 hours with complementary DNA of human Lrp4-green fluorescent protein, conferring green fluorescence to Lrp4-expressing cells. Membrane-bound anti-Lrp4 antibodies of positive controls were visualized by secondary antibodies exhibiting red fluorescence.

In the long exercise test, the patient, his uncle presenting with hypokalemic PP, and a male healthy control aged 30 years performed 7 rounds of finger spreading of 35 to 40 seconds each against strong resistance, followed by 2 to 3 seconds of relaxation, ensuring blood flow. The ulnar nerve was supramaximally stimulated at the wrist (constant current pulse, 0.2 milliseconds, 125%). Compound muscle action potentials were recorded from the abductor digiti minimi muscle immediately before and after exercise and then every minute for 5 minutes and finally every 5 minutes for up to 60 minutes.

The exons and exon-intron boundaries of *CACNA1S*,² *SCN4A*,³ *KCNJ2*,⁴ and *KCNJ18*⁶ were amplified from genomic DNA and bidirectionally sequenced using an automated 373A sequencer.

Results

Family History

The patient and his maternal uncle had hypokalemic PP with episodes of flaccid muscle weakness associated with low serum potassium levels and a pathologic decrement in the long exercise test (Figure 1). The spells were triggered by strenuous muscle work or exposure to a cold environment and improved by potassium intake. The patient's mother was deceased (stroke) and did not present with weakness episodes.


The Patient's History

The patient reported episodes of muscle weakness for 5 years. The episodes usually occurred during rest after strenuous exercise or after exposure to a cold environment and consisted of proximal leg weakness or tetraparesis. Bilateral stiffness of the thigh muscles occurred as abortive spells or followed the episodes of flaccid muscle weakness a few hours or a day later. Hypokalemia was detected during some episodes (lowest potassium value, 2.38 mEq/L [to convert to millimoles per liter, multiply by 1], whereas interictal serum potassium levels were in the normal range [3.5-5.5 mEq/L]). In his early 20s, attacks expanded from paraparesis to tetraparesis. During the first 2 years, the episodes disappeared spontaneously within 2 hours to a few days; however, in the last 3 years, potassium administration was required. During the last 6 months before referral, frequency and severity of attacks further increased and potassium administration became ineffective. The patient reported no other diseases.

Perioperative Findings

The patient was referred to 117 PLA Hospital with complete proximal tetraparesis and reduced muscle stretch reflexes. Although ocular, bulbar, and respiratory muscles seemed to be unaffected, a neostigmine test was performed and results were negative. Except for a reduced serum potassium level of 2.4 mEq/L, all other serum values, including

Figure 2. Images of the Enlarged Thymus



A, Mass without involvement of the surrounding tissue on the contrast-enhanced computed tomography scan of the anterior mediastinum. B, Hyperplastic thymus with lobular appearance and fibrous capsule. C, Lobular

architecture, cortex, medulla, and Hassall corpuscles and adipose tissue in the stroma (hematoxylin-eosin, original magnification $\times 100$). D, Immunostaining of cytokeratin 5/6 in hyperplastic epithelial cells (original magnification $\times 200$).

the autoantibodies listed in the Methods subsection, were not indicative of an autoimmune disorder. Particularly test results for the myasthenia-related autoantibodies, including anti-Lrp4 and the autoantibodies against various neuromuscular ion channels, were negative. Serum cortisol levels taken at 8 AM were in the normal range of 3.5 to 25.0 $\mu\text{g/dL}$ (to convert to nanomoles per liter, multiply by 27.588) (18.1 $\mu\text{g/dL}$ at admission and 15.9 $\mu\text{g/dL}$ at release) as well as serum ACTH level determined at release (24 ng/L; normal range, 10-90 ng/L). Thyroid and renal function, arterial blood pressure, electrocardiogram, radiography of the chest, and a contrast-enhanced CT scan of the adrenal glands revealed no abnormal findings. A contrast-enhanced CT scan of his chest exhibited an enlarged thymus (Figure 2A).

Thoracic surgery was performed right sided and video assisted under particular perioperative measures. In situ, an enlarged yellow-red thymus was observed. The excised thymus weighed 200 g (Figure 2B). Histologically, a lobular architecture with thymic cortex, medulla, Hassall corpuscles, and multiple hyperplastic follicles were observed, composed of adipose tissue and fibrous tissue (Figure 2C).

Immunohistology was negative for ACTH but revealed hyperplastic thymic epithelial cells positive for cytokeratin 5/6 (Figure 2D), confirming the diagnosis of follicular thymic hyperplasia.

Follow-up

Examinations performed 1 and 2 years after thymectomy revealed a postoperative absence of both abortive spells and weakness episodes. Genetic studies of both the patient and his uncle revealed no disease-causing mutation in the PP genes, and a contrast-enhanced CT scan of the uncle's chest was normal.

Additional Information on Unavailable Patients

We obtained the medical reports on (1) a male patient in his early 30s who had frequent hypokalemic weakness episodes and died of an acute paralytic attack associated with a serum potassium level of 2.0 mEq/L; thymic hyperplasia was found in the autopsy and (2) a female patient in her late 40s with familial hypokalemic PP and an enlarged thymus on the CT scan; thymectomy has not been performed because of the absence of end plate autoantibodies.

Discussion

Follicular thymic hyperplasia is seen in more than 50% of patients with myasthenia gravis. The remaining 50% have other autoimmune disorders, such as Graves disease or neuromyotonia, and tumors with increased ACTH or glucocorticoid levels.¹¹ Our patient exhibited normal ACTH, glucocorticoid, and thyroid serum levels. Additionally, we identified neither clinical hints for myasthenia gravis nor autoantibodies directed to the end plate. Finally, there were no autoantibodies against presynaptic calcium channels causing Lambert-Eaton syndrome or against dendrotoxin-sensitive potassium channels responsible for Morvan syndrome and neuromyotonia.¹²

In spite of negative screening results, the presence of autoantibodies in our patient cannot be excluded since even the majority of patients with asymptomatic thymic hyperplasia develop an autoimmune disorder.¹³ Evidence for contribution of a putative autoimmune disorder to the phenotype is the late onset and steady progression of the PP, suggesting a minimal required level of follicular thymic hyperplasia for clinical mani-

festation and a hyperplasia-proportional progression. The cessation of PP after thymectomy also supports this idea. Therefore, autoantibodies against sarcolemmal ion channels encoded by PP (susceptibility) genes should be tested in the future.

The reported familial recurrence indicates contribution of a hereditary disposition, probably due to a variant in an unknown susceptibility gene. This would explain the reduced penetrance in the mother and the lack of mutation in the PP channel proteins Cav1.1, Nav1.4, and Kir,²⁻⁴ despite the occurrence of hypokalemia-related weakness episodes, response to potassium administration, and abnormal long exercise test results.

We conclude that thymic hyperplasia can clinically manifest genetically predisposed hypokalemic PP. For patients with hypokalemic PP and late onset or clinical progression, we recommend screening for an autoimmune disease and, if thymic hyperplasia can be identified, thymectomy. Our findings may stimulate the identification of patients, raise the discussion on which patients should be screened and the need for thymectomy, and finally spark the search for autoantibodies against sarcolemmal ion channels.

ARTICLE INFORMATION

Accepted for Publication: June 17, 2013.

Published Online: September 23, 2013.
doi:10.1001/jamaneurol.2013.3918.

Author Contributions: Dr Lehmann-Horn had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: R. Yang, Wang, Lehmann-Horn.

Acquisition of data: R. Yang, Cao, Seelig, C. Yang, Liu, Pan, Zheng, Lehmann-Horn.

Analysis and interpretation of data: All authors.

Drafting of the manuscript: Cao, Wang.

Critical revision of the manuscript for important intellectual content: R. Yang, Jurkat-Rott, Wang, Seelig, C. Yang, Liu, Pan, Zheng, Lehmann-Horn.

Statistical analysis: R. Yang, Cao, C. Yang, Liu, Pan, Zheng.

Obtained funding: Lehmann-Horn.

Administrative, technical, or material support: R. Yang, Jurkat-Rott, Seelig.

Study supervision: R. Yang, Jurkat-Rott, Lehmann-Horn.

Conflict of Interest Disclosures: None reported.

Funding/Support: Drs Lehmann-Horn and Jurkat-Rott received grants from the nonprofit Else Kröner-Fresenius-Stiftung and the Federal Ministry of Education and Research (BMBF) for the IonNeuroNet rare diseases project. Dr Lehmann-Horn is an endowed Senior Research Professor of Neuroscience of the nonprofit Hertie Foundation.

Role of the Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Information: Correspondence on thymus surgery and reprint requests to Renrong Yang, MD, Department of Thoracic Surgery, 117 PLA Hospital, Hangzhou 310004, China (henryang88@gmail.com.cn), and correspondence on thymus pathology to Guofeng Wang, MD, Department of Pathology, Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou 310004, China (wgf2202@yahoo.com.cn).

Additional Contributions: Drs Lehmann-Horn and Jurkat-Rott thank Pavel Biletskiy, MD, and Henning Andersen, MD, for providing medical reports on 2 additional patients with hypokalemic PP and an enlarged thymus. We thank Brian Eyden, MD (Manchester, England), for improvements to the manuscript.

REFERENCES

- Platt D, Griggs R. Skeletal muscle channelopathies: new insights into the periodic paralyses and nondystrophic myotonias. *Curr Opin Neurol*. 2009;22(5):524-531.
- Jurkat-Rott K, Lehmann-Horn F, Elbaz A, et al. A calcium channel mutation causing hypokalemic periodic paralysis. *Hum Mol Genet*. 1994;3(8):1415-1419.
- Bulman DE, Scoggan KA, van Oene MD, et al. A novel sodium channel mutation in a family with hypokalemic periodic paralysis. *Neurology*. 1999;53(9):1932-1936.
- Plaster NM, Tawil R, Tristani-Firouzi M, et al. Mutations in Kir2.1 cause the developmental and episodic electrical phenotypes of Andersen's syndrome. *Cell*. 2001;105(4):511-519.
- Jurkat-Rott K, Weber MA, Fauler M, et al. K⁺-dependent paradoxical membrane depolarization and Na⁺ overload, major and

reversible contributors to weakness by ion channel leaks. *Proc Natl Acad Sci U S A*. 2009;106(10):4036-4041.

6. Ryan DP, da Silva MR, Soong TW, et al. Mutations in potassium channel Kir2.6 cause susceptibility to thyrotoxic hypokalemic periodic paralysis. *Cell*. 2010;140(1):88-98.

7. Cheng CJ, Lin SH, Lo YF, et al. Identification and functional characterization of Kir2.6 mutations associated with non-familial hypokalemic periodic paralysis. *J Biol Chem*. 2011;286(31):27425-27435.

8. Cheung CL, Lau KS, Ho AY, et al. Genome-wide association study identifies a susceptibility locus for thyrotoxic periodic paralysis at 17q24.3. *Nat Genet*. 2012;44(9):1026-1029.

9. Higuchi O, Hamuro J, Motomura M, Yamanashi Y. Autoantibodies to low-density lipoprotein receptor-related protein 4 in myasthenia gravis. *Ann Neurol*. 2011;69(2):418-422.

10. Zhang B, Tzartos JS, Belimezi M, et al. Autoantibodies to lipoprotein-related protein 4 in patients with double-seronegative myasthenia gravis. *Arch Neurol*. 2012;69(4):445-451.

11. Takahashi K, Al-Janabi NJ. Computed tomography and magnetic resonance imaging of mediastinal tumors. *J Magn Reson Imaging*. 2010;32(6):1325-1339.

12. Irani SR, Alexander S, Waters P, et al. Antibodies to Kv1 potassium channel-complex proteins leucine-rich, glioma inactivated 1 protein and contactin-associated protein-2 in limbic encephalitis, Morvan's syndrome and acquired neuromyotonia. *Brain*. 2010;133(9):2734-2748.

13. Singla S, Litzky LA, Kaiser LR, Shrager JB. Should asymptomatic enlarged thymus glands be resected? *J Thorac Cardiovasc Surg*. 2010;140(5):977-983.